

MEETING ABSTRACT

Open Access

Lynch II syndrome: a case report

B Scotto*, E Spera, G Merola, D Di Simone

From XXIII Annual Meeting of the Italian Society of Geriatric Surgery
 Lecce, Italy. 2-4 December 2010

Background

Lynch syndrome is an autosomal dominant cancer susceptibility syndrome that accounts for approximately 2–4% of all colorectal cancers (CRCs) and is caused by germline mutations of mismatch repair (MMR) genes [1-4]. It is characterized by an early onset of predominantly CRC and endometrial cancer (EC) as well as cancers of the stomach, small bowel, ureter and renal pelvis.

Materials and methods

We report a case of a patient 60 years old affected by synchronous primary cancers of the sigma and stomach diagnosed by TC. The patient had previously been operated on for the right hemicolectomy for adenocarcinoma, and hysterectomy for cancer. During hospitalization the patient has undergone surgery on the left hemicolectomy and a total gastrectomy. The patient underwent chemotherapy. Currently the patient is receiving chemotherapy for local recurrence.

Results

As a result of genetic study, the patient was found to be the first to have this mutation.

Conclusions

Surgery remains the front-line therapy for HNPCC. There is an ongoing controversy over the benefit of 5-fluorouracil-based adjuvant therapies for HNPCC-related colorectal tumors, particularly those in stages I and II. After reporting a null finding from their randomized controlled trial of aspirin (ASA) to prevent against the colorectal neoplasia of Lynch Syndrome, Burn and colleagues have recently reported new data, representing a longer follow-up period than reported in the initial NEJM paper. These new data demonstrate a

reduced incidence in Lynch Syndrome patients who were exposed to at least four years of high-dose aspirin, with a satisfactory risk profile. These results have been widely covered in the media; future studies will look at modifying (lowering) the dose (to reduce risk associated with the high dosage of ASA).

Published: 24 August 2011

References

1. Fishel R: The human mutator gene homolog MSH2 and its association with hereditary nonpolyposis colon cancer. *Cell* 1993, **75**:1027-38.
2. Bronner CE: Mutation in the DNA mismatch repair gene homologue hMLH1 is associated with hereditary non-polyposis colon cancer. *Nature* 1994, **368**:258-61.
3. Miyaki M: Germline mutation of MSH6 as the cause of hereditary nonpolyposis colorectal cancer. *Nat Genet* 1997, **17**:271-2.
4. Nicolaides NC: Papadopoulos Mutations of two PMS homologues in hereditary nonpolyposis colon cancer. *Nature* 1994, **371**:75-80.
5. Renda A, Relazione Biennale SIC: Multiple Primary Malignant. Springer-Verlag Italia; 2009.

doi:10.1186/1471-2318-11-S1-A58

Cite this article as: Scotto et al.: Lynch II syndrome: a case report. *BMC Geriatrics* 2011 **11**(Suppl 1):A58.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: segreteria@disalvo@gmail.com

Department of Surgical, Anesthesiologic and Emergency Sciences. University "Federico II", Naples, Italy